



# **SURVIVAL, PERSISTENCE, TRANSFER**

**An update on  
current knowledge  
on GMOs and  
the fate of their  
recombinant DNA**

by Beatrix Tappeser, Manuela Jäger,  
Claudia Eckelkamp

**TWN**

Third World Network

# **SURVIVAL, PERSISTENCE, TRANSFER**

*An update on current knowledge  
on GMOs and the fate of their recombinant DNA*

*Beatrix Tappeser,  
Manuela Jäger,  
Claudia Eckelkamp*

**TWN**

Third World Network  
Penang, Malaysia

***Survival, Persistence, Transfer — An update on current  
knowledge on GMOs and the fate of their recombinant DNA***

*is published by  
Third World Network  
228 Macalister Road  
10400 Penang, Malaysia.*

copyright © Third World Network 1999

This is part of a series of papers on biotechnology and biosafety that the Third World Network is publishing with a view to deepening public understanding on the ecological and safety aspects of the new biotechnologies, especially genetic engineering.

Printed by Jutaprint  
2 Solok Sungei Pinang 3, Sg. Pinang  
11600 Penang, Malaysia.

First Printing: 1999  
Second Printing: 2002

ISBN: 983-9747-37-1

# Contents

---

|                   |                                                                     |           |
|-------------------|---------------------------------------------------------------------|-----------|
| <b>Chapter 1.</b> | <b>Introduction</b>                                                 | <b>1</b>  |
| <b>Chapter 2.</b> | <b>Survival and Spread of GMOs</b>                                  | <b>3</b>  |
|                   | <i>Survival in the digestive tract</i>                              | 3         |
|                   | <i>Survival in wastewater and sludge</i>                            | 4         |
|                   | <i>Survival in aquatic ecosystems</i>                               | 5         |
|                   | <i>Survival in soil</i>                                             | 7         |
|                   | <i>Survival on plants</i>                                           | 11        |
| <b>Chapter 3.</b> | <b>Further Points to Consider: VNC state and adaptation ability</b> | <b>13</b> |
| <b>Chapter 4.</b> | <b>Spread of Cloned Sequences</b>                                   | <b>16</b> |
|                   | <i>Persistence of “naked” DNA</i>                                   | 17        |
|                   | <i>Persistence in wastewater treatment plants (water/sludge)</i>    | 17        |
|                   | <i>Persistence in aquatic systems</i>                               | 17        |
|                   | <i>Persistence in soils</i>                                         | 18        |
|                   | <i>Persistence in the digestive system</i>                          | 19        |
|                   | <i>“Avenues and barriers of genetic transmission” (Istok, 1991)</i> | 19        |
| <b>Chapter 5.</b> | <b>Discussion</b>                                                   | <b>24</b> |
|                   | References                                                          | 27        |

## Chapter 1

### Introduction

During the last years the release, whether tolerated or permitted, of genetically modified microorganisms (GMOs) and their nucleic acids into various environments has increased worldwide. Legislation concerning the containment of GMOs has been deregulated and safety measures have been relaxed throughout the industrial world, as genetic engineering has not occasioned any obvious accident or visible negative impact during the two decades of its rapid development and constantly increasing use. Another development of the last 10 to 15 years, according to the 1996 World Health Organisation (WHO) Report, has been an increase in frequency of outbreaks of new and reemerging infectious diseases. Moreover, current pathogen strains are often resistant to known treatments, some even to nearly all commonly used antibiotics. Horizontal gene transfer is now recognised to be the main avenue of exchange of genetic material in the microbial world, and hence also of the exchange and spread of antibiotic resistance genes.

These developments give rise to two questions. Does the extensive use of antibiotic resistance genes in genetic engineering contribute to the increase in frequency of antibiotic resistance in bacterial pathogens? And what will be the outcome of a spread of recombinant genes analogous to the spread of antibiotic resistance genes?

Do we have risk assessment procedures available for monitoring the fate of GMOs and their recombinant DNA? Are we capable of

detecting ecological impacts or health impacts in early warning systems? The answer has to be no, and this for various reasons. We are only beginning to understand microbial ecology. There is a lack of basic knowledge by which to judge the possible impacts of a given GMO or its recombinant DNA on different environments. In particular, we do not know enough of the special conditions under which gene transfer takes place. What are the selective conditions which facilitate the transfer of specific genes and may, for example, promote the transfer of recombinant gene constructs?

A prerequisite for the possibility of GMOs and recombinant DNA contributing to the spread of newly cloned genes is the viability and/or persistence of GMOs and their recombinant DNA in certain environments. Another point requiring analysis is the extent to which artificial vectors facilitate and/or enhance the probability of horizontal gene transfer.

In this review we present the latest data on survival of GMOs in different environments, persistence of recombinant DNA, and what is currently known of the different gene transfer mechanisms involved.

## Chapter 2

# Survival and Spread of GMOs

### Survival in the digestive tract

The digestive system of vertebrates and invertebrates alike offers possibilities for bacteria ingested with the food to get in close contact with each other and with the microflora of the animal. Its environment may prolong bacterial survival as compared with other environments and, due to its high density of microorganisms, increase the probability of gene transfer (Adamo and Gealt, 1996).

Moreover, bacteria can be disseminated by their new host animals, e.g. earthworms (Clegg et al., 1995). Some bacteria may even change their metabolic activities and capacity for survival during their passage through the digestive tract. (Clegg et al., 1995). Earthworm activities may also influence the rate of transfer of conjugative plasmids from *Pseudomonas fluorescens* to autochthonous soil microorganisms (Daane et al., 1996). It is also possible for transfer mechanisms other than conjugation to contribute to the dissemination of DNA from GMOs to the endogenous microflora of animals (Duval-Iflah, 1992).

Aside from these studies on bacterial interaction with the invertebrate digestive system, Brockmann et al. (1996) investigated the possibility of different strains of *Lactococcus lactis* colonising the rat intestine and the transfer of plasmids between these bacteria. While the digestive tract of germfree animals was found to be rapidly colonisable by all kinds of bacteria, this only succeeded temporar-

ily (for 2 - 3 days) in non-germfree controls. Studies with *Lactococcus lactis* in the human intestine yielded similar results (Klijn et al., 1995a). Moreover, the conjugative plasmid pAM $\beta$ 1, which has a broad host-range, proved transferable both to a *Lactococcus* recipient strain in the intestine of germ-free rats and an endogenous strain of *Enterococcus faecalis* in that of non-germfree animals (Brockmann et al., 1996). These authors conclude that even those microorganisms only temporarily detectable in the digestive tract are able to transfer their conjugative plasmids to the endogenous microflora. The rate of such events mainly depends on the structure of the plasmids themselves.

## **Survival in wastewater and sludge**

The most important factor influencing bacterial survival in wastewater has to do with competitors. Because predator populations fluctuate seasonally, the time of year is an important factor in evaluating the risk of GMOs' survival (Inamori et al., 1992). If they are not eliminated by their competitors, *E. coli* K12 and *Pseudomonas putida* species have the best chances of survival when associated with snow particles containing sludge material (McClure et al., 1989; Overbeck, 1991). Whereas viable cell counts in supernatant were found to decrease quickly after introduction of recombinant *E. coli* K12 into the aeration basin of a model wastewater treatment plant, this was not usually the case in the remaining bacteria attached to sludge particles (Heitkamp et al., 1993).

Most studies on survival and gene transfer in wastewater and sludge use *in vitro* systems or models of settlement tanks or activated sludge for investigation. Ashelford et al. (1995) concentrated on percolating filter beds, which are layered with living cells (a so-called biofilm) similar to the river epilithon. Two inoculated *Pseudomonas* strains survived the whole investigation time of 145 days. The authors demonstrated the transfer of a conjugative plasmid (pQKH6) harboured by one of these strains to another,



initially plasmid-less, strain (Ashelford et al., 1995).

Similar results were also published by Feldmann and Sahm (1994), who investigated the survival of four different recombinant microorganisms in laboratory wastewater systems.<sup>1</sup> While none of the GMOs persisted in the settlement tank, two yeasts tested in the aeration basin remained detectable for more than 25 days.

## Survival in aquatic ecosystems

A lot of the above data apply in a like manner to aquatic ecosystems. The chance of survival of, e.g., *E. coli* and *Campylobacter jejuni* increases in filtered water taken from lakes, since filtering eliminates predators and competitors (Korhonen and Martikainen, 1991). Sterile tapwater also allowed the persistence of four transgenic microorganisms tested within the scope of the "Verbundprojekt Sicherheitsforschung Gentechnik"<sup>2</sup>: in this medium survival lasted the whole testing period of 440 days (Tebbe et al., 1994a).

Brettar et al. (1994) collected data on the survival of *Pseudomonas putida* DSM3931 in lakewater mesocosms. Strains were not genetically modified and survived the whole testing period of 10 weeks. Mesocosms additionally inoculated with culture media even induced an increase of bacterial numbers during the first 10 days. The most important factor responsible for the reduction of bacterial populations was predation, though many bacteria escaped by associating with particles and/or sediments during the first two days (Brettar et al., 1994).

Predation through protozoa also eliminated most bacteria

---

<sup>1</sup> These experiments were part of the "Verbundprojekt Sicherheitsforschung Gentechnik", which aims at investigating the safety of GMOs released from production facilities, and were sponsored by Bayer AG, Germany.

<sup>2</sup> *Saccharomyces cerevisiae*, *Zymomonas mobilis*, *Corynebacterium glutamicum*, and *Hansenula polymorpha*.

(*Pseudomonas fluorescens* AG1) inoculated in seawater mesocosms, though the extent of this was found to depend on the state of bacterial growth (Christoffersen et al., 1995). Elimination of bacteria in seawater is also achieved by lysis through bacteriophages (bacterial viruses), depending on seasonal changes (Weinbauer et al., 1995).

Survival of bacteria in aquatic ecosystems also depends on abiotic factors like temperature, nutrient availability and osmotic conditions (Ahl et al., 1995). These authors studied the survival of *Pseudomonas fluorescens* Ag1 in experiments with seawater microcosms (0.5 l water) and seawater mesocosms (5,300 l water). Ahl et al. (1995) demonstrated a less pronounced reduction of bacterial cells inoculated in mesocosms as compared with earlier microcosmic experiments. Of the inoculated *Pseudomonads* 25% survived the whole testing period of 14 days.

Intensity and seasonal changes of light are further important abiotic factors influencing the survival of bacteria in natural environments (Canteras et al., 1995). Five different strains of *E. coli* kept in complete darkness persisted for 96 hours in sterile riverwater with only a minor reduction of cfu<sup>3</sup>. Differences in persistence were only apparent upon illumination. However, sterile conditions cannot be taken to represent natural aquatic ecosystems. In the comparative study of Alvarez et al. (1996b) on the survival of *E. coli* strains with or without plasmids (DH1 and JM103) in tropical riverwater systems, differences in survival only showed in unsterile test-chambers.

Bacteria taken from their natural environment, transformed with a plasmid and then released again as GMOs, have a very high chance of establishing themselves in their original ecosystem. For example, Sobecky et al. (1996) demonstrated that a marine bacterium (*Achromobacter* sp.) is able to form stable populations in seawater

---

<sup>3</sup> cfu = colony-forming units

microcosms within 2-3 weeks after being transformed with a plasmid bearing a gene encoding an alkaline phosphatase. Even the enzyme was proved to have retained its activity.

Bacteria not adapted to environmental conditions, like typical enteropathogenic microorganisms (e.g. *E. coli* ETEC, *Yersinia enterocolitica* and *Campylobacter jejuni*) are not only able to survive within their hosts at body temperature but may also persist in a cultivable stage for weeks in aquatic environments at temperatures between 6°C and 16°C (Terzieva and McFeters, 1991). Davies et al. (1995) demonstrated that the time of survival of enteropathogenic bacteria (faecal *Streptococci* and Coliforme) in sediments exceeds that in open water. While gram-negative bacteria inoculated in open seawater soon entered a stage of non-culturability, *E. coli* cells residing in sediments remained cultivable during the whole testing time of 68 days (Davies et al., 1995).

The same seems to be true of groundwater aquifers and freshwater environments, where sediments provide a better chance of survival than open water. This was demonstrated by Winkler et al. (1995) with *Burkholderia cepacia* in a groundwater microcosm experiment and by Fish and Pettibone (1995) with *E. coli*. The latter species remained cultivable in sediments for 56 days, i.e., for the entire duration of the test.

## Survival in soil

Since soil environments are especially complex and heterogeneous systems, it is rather difficult to predict the chance of survival they offer to newly introduced GMOs. **The density of soil bacteria is estimated at about 108 to 109 cfu/g soil. However, less than 10% of these are cultivable using current techniques, and only around 1% have been characterised to date (Hugenholz and Pace, 1996; Pace, 1997).** In clay sediments bacteria are detectable at depths down to 224 m (Boivin-Jahns et al., 1996). Microbial populations from the laboratory decrease drastically during the first days after

their introduction into natural soil, but then often remain at a constant level (Schmidt, 1991).

It is most important in assessing and comparing published data on the survival of GMOs in soil systems to thoroughly analyse the design of the microcosm or mesocosm and detection method being used. The variety of microcosms studied makes it is rather difficult to directly compare the results collected (Angle et al., 1995). An undisturbed soil texture is an important prerequisite for optimal imitation of natural environmental conditions, as Angle et al. (1995) demonstrated. These authors compared survival of *Pseudomonas aureofaciens* in microcosms and deliberate release experiments in naked soil for two years. Validation of the microcosms used is also expedient with aquatic systems. Leser (1995) found 63%-76% agreement between data obtained from a lakewater microcosm and a natural lake. After inoculation of *Alcaligenes eutrophus* in both systems the degree of coincidence was reduced to 2-27%, but increased again with decreasing cell counts (Leser, 1995).

As with aquatic and wastewater systems, in soils, too, predation through protozoa is one of the most important factors influencing the persistence of inoculated GMOs. In these environments protozoal activity requires a certain degree of soil moisture and free water in soil pores of suitable size (Van Elsas 1992; Wright et al., 1995). Experiments in soil microcosms with *Pseudomonas fluorescens* and the ciliate *Colopoda steinii* demonstrated that small pores (< 6µm) may offer effective protection zones against predators (Wright et al., 1995). While bacterial populations in larger pores (< 30µm) diminish more rapidly, bacteria that do survive show a higher level of metabolic activity in response to the richer supply of nutrients there (Wright et al., 1995). Van Der Hoeven et al. (1996) used a computer simulation model to evaluate the chance of survival of GMOs in soil pores of different sizes. Bacteria introduced into the soil were eliminated very slowly, in some cases taking years for total elimination (Van Der Hoeven et al., 1996).

The number of bacteria inoculated into the soil is a very important determinant of the time of possible survival, as differences as small as 100 *E. coli* K12 cells/g soil suffice to change the results of an assay (Recorbet et al., 1992). This strong correlation between inoculation number and survival is also found in aquatic ecosystems (Karapinar and Sahika, 1991). Another criterion is the depth of inoculation. In the range from 10 to 50 cm the chance of survival of the phytopathogenic bacterium *Erwinia carotovora* increases with growing inoculation depth. This correlation largely reflects differences in competitive pressure (Armon et al., 1995). Recorbet et al. (1995) demonstrated uneven distribution of genetically modified *E. coli* which had been inoculated in soil systems. Furthermore, GMOs were concentrated in different soil layers than autochthonous bacteria.

Spore-forming bacteria like *Bacillus subtilis* are particularly likely to persist for prolonged time periods. Spores of *Bacillus subtilis* of both native and genetically modified parent strains remained culturable in sterile soil for at least 50 days (Tokuda et al., 1995). GMOs' ability to adapt to natural environments makes it more difficult to estimate their chance of survival after release. Cells of *Pseudomonas fluorescens* showed improved stress resistance against temperature or osmotic pressure one day after inoculation into two different types of soil (Van Overbeek et al., 1995). When cultivated under starving conditions in the laboratory these bacteria may stand stress-factors like "hunger" for 70 days by reducing their cell size. These physiological changes are soon reverted again when conditions are made to improve (Clegg et al., 1996).

There is still no evidence of plasmid-bearing microorganisms being less fit than their native parents: e.g. Fujimura et al. (1994) tested genetically modified *Saccharomyces cerevisiae* and their parent strains under different environmental conditions but were unable to prove any negative impact in terms of survival probability. Under selective pressure, exerted, for example, by antibiotic substances, GMOs

may even profit from possessing the corresponding resistance genes (Van Elsas, 1992). This was also demonstrated by Ramos et al. (1994) with recombinant *Pseudomonas putida* (pWW0-EB62) in soil systems. In unsterile soil and under assumed selective pressure cells numbers even increased after inoculation (Ramos et al., 1994). However, application of selective pressure is not a necessary prerequisite for survival of GMOs.

Sjogren (1995) initiated a very profound project on survival of *E. coli* using a multiresistant native strain isolated from a lake. The bacteria were inoculated into different field plots seeded with rye. There were no more bacteria detectable on the rye grass 41 days later. After two months bacteria were reisolated from soil at a depth of 20 to 50 cm. At 60-cm depth they reached groundwater and persisted there at the water-soil interface for two more years. Six years later 1.5 cfu/ml were still detectable, and after eight years cell counts had increased to 8 cfu/ml. After 13 days, at the end of the experiment, population density was still at 0.25 cfu/ml (Sjogren, 1995). The author calculated the speed of distribution as about 2 cm/day. In analogous experiments performed under laboratory conditions with microcosms of the same soil the maximum survival time had been 2.5 years (Sjogren, 1995).

GMOs intended for use in natural environments are designed to persist and show at least some competitive ability versus autochthonous microorganisms. Experiments with *Rhizobium meliloti*, which had been genetically modified for biological containment by disrupting a gene for recombination, only demonstrated reduced survival of the genetically modified as compared with the native strain when performed in model ecosystems, but not after deliberate release in agricultural soils (Dresing et al., 1995; Selbitschka et al., 1994).

Beside surviving, GMOs inoculated in soil may also be distributed further, thus increasing the chances of gene transfer. Gillespie et al. (1995) simulated rainfall on wheat plants or naked soil in green-

house experiments after inoculation with genetically modified *Pseudomonas aureofaciens*. The number of cells detectable in the soil was found to correlate with the number of cells in water running from the plots. One exception to this was that 245 days after inoculation 108 cfu were measured in the water probes, whereas cell counts in the soil were below the detection limit. The authors concluded that standard techniques for detection of GMOs in soil may not be suitable for registering bacteria in all soil layers and that short and heavy rainfalls can cause quite high numbers of GMOs (about 1,014 cfu/ha) to be washed away from the place of release (Gillespie et al., 1995). Similar testing was done by Hekman et al. (1995) with *Pseudomonas fluorescens* and *Burkholderia cepacia*. In the case of planted microcosms, these authors demonstrated enhanced distribution of inoculated bacteria in deeper soil layers after moderate rainfall (Hekman et al., 1995).

## Survival on plants

It is evident from the last paragraph that any analysis of survival of GMOs in soil has to include the plants growing there. A recombinant strain of *Bacillus amyloliquefaciens*, used to produce (-amylase) was no longer detectable in the soil of planted microcosms six days after inoculation, though it persisted on the grass for at least 70 days (Wendt-Potthoff et al., 1994). Genetically modified strains of the phytopathogenic bacteria *Xanthomonas campestris* inoculated on cabbage host plants survived for six months and were distributed to other cruciferous plants and even unrelated plants (Dane and Shaw, 1996).

Persistence of bacteria inoculated on plants may depend on the plant species used. The population density of two strains of *Pseudomonas fluorescens* remained at a constant level on leaf surfaces of capsicum and eggplants during a 30-day glasshouse experiment, but decreased on leaves of strawberry and tomato (Cirvilleri et al., 1996). Genetically modified *Pseudomonas fluorescens* whose parent strain had been isolated from sugar beets persisted

on leaf surfaces of sugar beets in the glasshouse for at least 531 days. Fluctuations in the number of the inoculated bacteria correlated with fluctuations in the endemic *Pseudomonads*. Besides sometimes being the most prominent population on the leaves, GMOs proved capable of colonising other plant species in the glasshouse and of further distribution by insects (Thompson et al., 1995a). Analogous deliberate release experiments demonstrated survival of these GMOs for 270 days (the whole vegetation period) and confirmed their capability of spreading further (Thompson et al., 1995b). Even sugar beets planted in the following season were colonised, though cell counts fell below the detection limit (1-20 cfu/g) during the winter season in both soil and plant tissue (Thompson et al., 1995b).

De Leij et al. (1995), too, have studied the survival and dispersal of *Pseudomonas fluorescens* on wheat plants. These authors demonstrated that the strain is able to survive longer in field experiments than in microcosm or greenhouse studies and can disperse to other plants and the soil (De Leij et al., 1995). Variation in survival time in different field experiments is due to the variety of soil types used. This is confirmed by Kinkel et al. (1996) whose data from microcosm experiments on the persistence of *Pseudomonas syringae* on different plant species did not correlate with data from release experiments. Bacteria inoculated in field testing produced smaller or larger populations than those subjected to laboratory testing, depending on which plant species was being used.



## Chapter 3

### Further Points to Consider: VNC state and adaptation ability

**A** clear distinction must be made between cultivable and dormant cells, regardless of the environment under study (Dott et al., 1991; Turpin et al., 1993). The dormancy problem is currently receiving more and more attention, since results have been found to differ tremendously according to whether all cells or only cultivable cells (Kragelund and Nybroe, 1996; Pickup, 1991; Stotzky et al., 1991) are counted.

These VNC, i.e., viable non-culturable, bacteria are predominant in marine environments with high salt concentration (Byrd and Colwell, 1990; Garcia-Lara et al., 1993). There are various especially adapted methods of detecting VNC cells and cultivable cells. However, counts of VNC bacteria in environmental probes are usually rather variable (Bianchi and Giuliano, 1996; Byrd and Colwell, 1990; Jensen et al., 1996; Zweifel and Hagström, 1995). The problem of poor culturability has also to be addressed in the case of bacteria in activated sludge or soil probes (Binnerup et al., 1993; Wagner et al., 1993).

VNC cells can be distinguished from dead cells by some metabolism and protein synthesis parameters. Moreover, rather than lose their plasmids, they may even enrich their intracellular plasmid content (Arturo-Schaan et al., 1996; Nwoguh et al., 1995; Nybroe et al., 1996). These so-called dormant cells can be activated again when environmental conditions change. This is not to say that an im-

proved nutritional state is the only factor responsible for reactivation; temperature fluctuations in natural environments may also play a role (Colwell et al., 1996; Ferguson et al., 1995; Jiang and Chai, 1996; Oliver et al., 1995a; Rahman et al., 1996). The presence of plasmids can further contribute to induction of the VNC stage (Oliver et al., 1995b). Bacteria detection methods which rely on the expression of marker genes and are thus blind to VNC cells should therefore at least be combined with techniques that record VNC cells (Lee et al., 1996; Leff and Leff, 1996; Oliver et al., 1995b).

Obviously, the choice of a detection method has a great influence on the record of bacteria released in natural environments. The ability of microorganisms to adsorb to particles and sediments renders detection more difficult, though application of PCR has certainly lowered detection limits (Bej et al., 1990; Hodson et al., 1995; Lindahl 1996; Lindahl and Bakken, 1995; Tsushima et al., 1995; Vahjen and Tebbe, 1994). One disadvantage of this technique is the remaining difficulty in distinguishing between isolated DNA and DNA associated with living or VNC organisms (Brockmann et al., 1996; England et al., 1995; Tamanai-Shacoori et al., 1996).

A further point to consider in the context of survival of GMOs in natural environments is their capability to adapt to unfavourable environmental conditions. When given a chance to adapt slowly, bacterial populations in aquatic microcosms imitating natural surroundings show prolonged persistence as compared to non-adapted populations of the same strain (Awong et al., 1990; Mezrioui et al., 1994; Sobecky et al., 1992). Bacteria cultivated in the laboratory while exposed to various primary stress factors like unfavourable pH or nutrient deficiency are able to adapt, and also show higher resistance against different secondary stress factors (Ferianc et al., 1995; Lou and Yousef, 1996). Consequently, any design of microcosm or mesocosm experiments should include at least some environmental stress factors in order to check GMOs' responsiveness to them. In determining the time-scale of such experiments one has to consider that adaptation may be a slow process and that populations

of GMOs no longer detectable after a short time may multiply and rise again (Clegg et al., 1995; Gillespie et al., 1995; Sjorgen, 1995; Ramos et al., 1994; Thompson et al., 1995b). Stress factors may even act as a positive selection pressure.

## Chapter 4

# Spread of Cloned Sequences

Risk assessment should not be satisfied with the assumption or proof that a given GMO will not survive. It must rather extend to the fate of the DNA, which may be stably integrated, eventually expressed and may by chance provide advantages for indigenous microorganisms (Henschke and Schmidt, 1990; Istock, 1991; Schmidt et al., 1994). GMOs may only require a short time of survival in order to be transported to other ecological niches or be able to transfer some of their nucleic acids to other members of a given ecosystem (Byzov et al., 1993; Heijnen and Marinissen, 1995; Weiskel et al., 1996). While conjugation and transduction are restricted to living cells, transformation may also take place after the death of donor cells. The probability of gene transfer rises with the stability and quantity of isolated DNA (Fulthorpe and Wyndham, 1991), though selective forces have also to be taken into account and may play a crucial role.

For example, Pukall et al. (1996) isolated plasmids with mobilising capability from soil probes and pig manure in numbers substantially higher than estimated. Such plasmids can transform competent cells living in soil or manure environments also under suboptimal conditions and may even be able to mobilise recombinant plasmids in these cells (Lorenz and Wackernagel, 1987; Pukall et al., 1996).

## **Persistence of “naked” DNA**

Isolated DNA does not only originate from dead and lysed cells but can also be actively secreted (Paget and Simonet, 1994). Quantifying extracellular DNA in environmental media seems to pose difficulties, judging by the great variability of published data (Lorenz and Wackernagel, 1994). Some authors assume that bacteria in natural environments release large quantities of high molecular DNA and plasmids into the extracellular gene pool (Lorenz and Wackernagel, 1994; Romanowski et al., 1990; Wackernagel et al. 1992). Secretion of DNA and competence for nucleic acid uptake are induced in particular in unfavourable environmental conditions like starvation (Lorenz and Wackernagel, 1994; Wackernagel et al., 1992).

## **Persistence in wastewater treatment plants (water/sludge)**

Wastewater treatment plants guarantee rapid and almost complete inactivation and degradation of isolated DNA when it is suspended in open water. Nucleic acids adsorbed to sludge particles, however, find some degree of protection against degradation and also against detection (Gross et al., 1994; Aardema et al., 1983; Bauda et al., 1995). As efficiency of inactivation of naked DNA depends on different factors like temperature, it varies with the season and type of treatment plant under study (Bergemann et al., 1994; Lorenz and Wackernagel, 1994).

## **Persistence in aquatic systems**

Some recent studies have dealt with the origin and distribution of naked DNA in aquatic systems (Beebee 1993; Hermansson and Linberg, 1994; Jiang and Paul, 1995; Weinbauber et al., 1995). As in wastewater environments, isolated and dissolved nucleic acids are

rapidly degraded in aquatic systems by enzymatic activity (Alvarez et al., 1996a; Lorenz and Wackernagel, 1994). Adsorption to sand or clay minerals protects DNA against nucleases even if these are bound to the same particles (Aardema et al., 1983; Romanowski et al., 1993a).

Prolonged protection of nucleic acids against inactivation is also provided by VNC cells (Byrd et al., 1992; Lorenz and Wackernagel, 1994). If naked DNA remains biologically active, this can give rise to the transformation of competent cells. Therefore, transformation experiments are a suitable method of detecting intact molecules, regardless of whether they were adsorbed to particles or not (Alvarez et al., 1996a; Chamier et al., 1993; Mieschendahl and Danneberg, 1994; Romanowski et al., 1993a).

## **Persistence in soils**

Isolated DNA in non-sterile soils can persist for some time, depending on various factors like temperature, pH, salt concentration, type of soil and nucleic acid material, all of which may influence the binding of DNA on mineral or quartz particles (Lorenz and Wackernagel, 1992). As in aquatic systems, when introduced into non-sterile soils, particle-adsorbed nucleic acids are protected against degradation by nucleases, even if these are bound to the same particle samples (Khanna and Stotzky, 1992; Lorenz and Wackernagel, 1987). Nucleic acids can bind to minerals quite rapidly and relatively independently of pH (Lorenz and Wackernagel, 1994). One factor determining the speed of association in aquatic systems is the structure of the DNA (Khanna and Stotzky 1992; Lorenz and Wackernagel, 1992; Lorenz et al., 1988; Paget et al., 1993). Factors relevant to soil environment include the humidity and granular structure of the soil (Hobom, 1995).

Data on the persistence of isolated DNA in soils show great variation. Smalla (1995) detected recombinant DNA from sugar beets in soil 18 months after inoculation of plant material. Romanowski et

al. (1993b) found that particle-associated plasmids could persist in different soils for at least 60 days. Plasmid conformation remained intact during the first two days in this study. According to Wackernagel et al. (1992) and Romanowski et al. (1992), the time-limit for detecting plasmids in unsterile soil probes without PCR application is five to ten days. Intact genes or plasmids can frequently be detected for as long as the originating bacteria.

### **Persistence in the digestive system**

Consequently, if isolated DNA is able to persist in different environmental media, then it is quite possible for the digestive system of animals or humans to take up these nucleic acids with the food or drinking water. Moreover, as Schubbert et al. (1996) demonstrated in feeding experiments on mice, nucleic acids reaching the gastrointestinal tract must not necessarily be completely fragmented, and hence inactivated, but may just as well reach the bloodstream and temporarily even be detected in leukocytes and liver cells. The transferability of these results from mice experiments to other mammalian digestive systems, including the human one, is an obvious possibility. Investigating the persistence of bacterial cells (*Lactococcus lactis*) and their DNA in the human intestine Klijn et al. (1995b) were able to isolate the bacterial DNA from faeces four days after administering the inoculated milk drinks.

### **“Avenues and barriers of genetic transmission” (Istock, 1991)**

To date gene transfer has mostly been found by means of specially designed plasmids, as in a number of studies carried out in microcosms (Collard et al., 1993; Del Solar et al., 1993; Mazodier and Davies 1991; Miller et al., 1992; Paget et al., 1992; Paul et al., 1991; Paul, 1992). However, the spread of antibiotic resistance markers throughout bacterial communities shows that gene transfer is likely to happen not only in more or less artificial settings but also under natural conditions (Saye and Miller 1989; Götz et al., 1996; Kruse

and Sorum, 1994). Moreover, some plasmids originating from gram-positive bacteria have also been isolated in gram-negative bacteria, stressing the possibility of a wide distribution of genetic information. It was not possible to ascertain in this case whether gene transfer took place via conjugation, transformation or transduction (Del Solar et al., 1993).

**Conjugation**, i.e., the transfer of nucleic acids by direct contact between bacterial donor and recipient cells, is assumed to be the most effective mechanism of genetic transmission under natural conditions. Wastewater treatment plants offer perfect conditions for this kind of transmission because of their abundance of cell populations (Schneider et al., 1994). Analogously, conjugation in the gastrointestinal tract is promoted by high population densities of potential bacterial donor and receptor cells (Morrison, 1996). Several recent studies have dealt with conjugation in digestive systems (Adamo and Gealt, 1996; Brockmann et al., 1996; Klijn et al., 1995a,b; Rang et al., 1996; Rybachenko et al., 1996). Microcosm experiments or *in situ* testing with soil or aquatic systems also demonstrate the transfer of plasmids by conjugation in other environments (Lebaron et al., 1994; Sandaa and Enger, 1994; Bale et al., 1988; Pukall et al., 1996; Lukin and Prozorov, 1992). Barkay et al. (1995) have pointed out the importance of using autochthonous bacteria as donors or recipients for efficient detection of transfer through conjugation in natural environments. Other authors have discussed the influence of different soil systems and supplements of plant material, temperature and soil humidity (Ashelford et al., 1995; Hermansson and Linberg, 1994; Klingmüller, 1993; Temann et al., 1992; Van Elsas et al., 1990). The main factors inhibiting or promoting conjugation in natural environments are thought to be selective forces from, e.g., heavy metals, herbicides, or antibiotics (De Rore et al., 1994a,b; Gadkari, 1991; Klingmüller and Rieder, 1994; Kozdroj and Pietrowska-Seget, 1995). Lafuente et al. (1996) have pointed out that conjugation in sterile soil microcosms is most efficient in "natural" abiotic conditions. As conjugation is an energy-consuming process, it can be promoted by any of a variety of



nutrients (Götz and Smalla, 1997). Any environmental compartment fulfilling this requirement like sediments, biofilms or water-air interfaces, and additionally characterised by high numbers of bacterial cells, will provide optimal conditions for conjugation (Amabile-Cuevas and Chicurel, 1996; Barkay et al., 1995; Hermansson and Linberg, 1994).

**Transduction**, a gene transmission process mediated by bacteriophages, requires living cells. It is more likely to occur under conditions of high metabolic activity (Ripp and Miller, 1995). Transduction may contribute to the spread of plasmids which are not transferable by conjugation for lack of essential sequences (Replicon et al., 1995). It is not restricted to plasmids, as chromosomal DNA can be transferred as well (Lorenz and Wackernagel, 1993). In this case the host range of the phages determines the possible radius of distribution of genetic material. Empirical evidence of gene transfer by transduction also taking place in natural environments (Ogunseitan et al., 1992; Ripp et al., 1994; Schicklmaier and Schmieger, 1995; Stotzky and Babich, 1994) has been further substantiated by experiments carried out under natural conditions (Kidambi et al., 1994; Miller et al., 1992) and by the detection of high numbers of phage particles in some environmental media (Kokjohn and Miller, 1992; Lorenz and Wackernagel, 1993; Schäfer, 1996). As transduction requires a certain minimum concentration of phage particles and corresponding bacterial host cells in order to be efficient, it is more likely to occur in environmental compartments characterised by high cell density and plenteous nutrients (Ogunseitan et al., 1992; Replicon et al., 1995). This is not an unconditional prerequisite for transduction, however, since also oligotrophic conditions and natural cell densities allow transduction (Goodman, 1994; Ripp et al., 1994). Adsorption of phages and bacteria to particles further increases the chances of survival and interaction and, consequently, also of transduction (Pickup et al., 1993; Ripp and Miller, 1995).

**Transformation**, the third mechanism of gene transfer, does not

depend on living cells as donors of nucleic acids. It requires the active uptake of isolated DNA by recipient cells which are in a physiological stage of competence (Lorenz and Wackernagel, 1994). According to present knowledge, induction of competence is determined by growth stage, impact of stress factors and/or presence of diffusible competence factors secreted by bacteria, and, not least, by the species in question (Baur et al., 1996; Cheng et al., 1997; Goodman, 1994; Lorenz and Wackernagel, 1994; Prozorov, 1997; Schlüter and Potrykus, 1996). The next step towards establishment of the foreign nucleic acids involves recombination as a means of integrating the new sequences into the recipient genome. Homologous sequences have better chances of transformation, though uptake does not usually depend on the presence of similar DNA (Lorenz and Wackernagel, 1994; Prozorov, 1997).

Establishment of plasmids does not depend on homology, **but only on suitable origins of replication** (Lorenz and Wackernagel, 1994). Experimental and empirical data both suggest that transformation is common in natural environments (Boyle-Vavra and Seifert, 1996; Hermansson and Linberg, 1994; Lorenz and Wackernagel, 1994; Schäfer, 1996). Microorganisms in aquatic and soil systems can also be transformed by isolated DNA, as has been demonstrated by Frischer et al. (1994), Lorenz and Wackernagel (1994), Paget and Simonet (1993) and others. Transformation of endogenous bacteria in the gastrointestinal tract was shown in the intestine of *Folsomia candida* after feeding with GMOs (Hobom 1995; Tebbe et al., 1994b). Baur et al. (1996) postulate that natural environments can provide optimal conditions for transformation which are not properly imitable in the laboratory. Environmental factors influencing transformation include lack or oversupply of nutrients and certain minerals, ionic strength of water and temperature (Baur et al., 1996; Frischer et al., 1993; Hermansson and Linberg, 1994; Lunsford and London, 1996; Williams et al., 1996). As in the case of conjugation and transduction, enhanced rates of transformation are found on surfaces and biofilms (Baur et al., 1996; Hermansson and Linberg, 1994).

Reports of intense natural gene transfer may appear to stand in contradiction to the ability of bacterial strains and varieties to maintain their genetic make-up. However, there are restriction systems whose action consists much rather in degrading or “silencing” foreign DNA than in preventing uptake of foreign DNA (Heinemann, 1991). Plasmid incompatibility is another means of hindering genetic transfer (Naik et al., 1994). Stressful conditions seem to reduce the activity of restriction systems in bacteria and to induce genes mediating recombination (Saunders and Saunders, 1993; Schäfer et al., 1994).

On the one hand, special plasmids are being designed to reinforce the constraints of biological containment (safety vectors without transfer genes), while on the other, progress in cloning relies on shuttle vectors capable of transgressing existing borders between bacterial classes and even kingdoms (Doucet-Populaire, 1992; Henchke and Schmidt 1990; Schäfer et al., 1994; Trieu-Cuot et al., 1987). Plasmids are being constructed to undermine the different levels of restriction they encounter under normal conditions in a natural environment and so ensure their persistence and integration (Dunn et al., 1993; Heinemann, 1991).

## Chapter 5

### Discussion

Current security measures for research and production are based on earlier assumptions about the survival and transfer abilities of microorganisms. In our opinion, the data in this article expose the concept of biological containment, on which legal regulation is based, as unsound, and this for several reasons:

- GMOs can survive or transfer their transgenes to indigenous organisms;
- DNA is more stable than has been hitherto imagined;
- and DNA taken up with the food is not completely degraded in the gastrointestinal tract, but has rather been found to enter white blood cells and spleen and liver cells.
- DNA can even be transferred to the cells of foetuses, as has been shown in newborn mice. Here transfer probably took place via the placenta (Doerfler and Schubbert, 1997).

Moreover, there are now nucleic acid constructs containing sequences which contribute not only to effective replication in different cellular backgrounds but also to stability and integration via recombination, transfer and extraordinary expression. They are — this is the essence of the art — especially designed to fulfil these jobs.

Moreover, neither laboratory nor *in situ* studies on survival, DNA persistence and gene transfer provide a realistic measure of what can happen in complex terrestrial, aquatic and gut environments (Spielman et al., 1996; Stotzky et al., 1991). Crude as they may be, validated microcosm and mesocosm studies imitating natural conditions as accurately as possible are nevertheless an indispensable means of testing and prognosticating the influence of some of the biotic and abiotic factors operating on GMOs in natural environments (Pickup et al., 1993; Spielman et al., 1996).

What has been neglected until now in designing such studies are the selective forces which may act on the spread of recombinant genes. Especially in disturbed or polluted environments, contaminants like heavy metals or high salt concentrations may possibly facilitate gene transfer.

In a recent seminar organised by the Norwegian Biotechnology Advisory Board on Antibiotic Resistance, Marker Genes and Transgenic Plants one of the invited speakers summarised:

"The extent and consequences of horizontal gene transfer are apparent in the evolution of antibiotic resistant microorganisms and evidence suggests that horizontal gene transfer may be equally frequent among multicellular eukaryotic organisms. But the actual and potential frequencies of gene transfer are poor indicators of risk; very common genes are not maintained in nature if unselected and rare genes become common extremely quickly if they are the subject of selection. What remains essential to assessing risk is identifying all potential selective pressures a recombinant gene might be suited to neutralise. New evidence suggests that current knowledge of evolutionary theory is inadequate to predict the fate of recombinant organisms or recombinant genes." (Heinemann, 1997).

To this we would add that until there is sufficient scientific evidence that a given GMO or its recombinant genes will not pose

any environmental stress or health impact, we should abide by the precautionary approach in regulating the contained use and deliberate release of GMOs. This implies putting a stop to deregulation in favour of contained use and to tolerated releases from production plants where environmental impacts are not routinely assessed. We should bear in mind that soil ecology, for example, has much to do with mineralisation and nutrient flow, which again is dependent on the enzymatic properties of the organismal network. Genes currently perceived as harmless like many enzyme-coding genes (or the organisms containing them) may alter soil chemistry and so create selective pressure or conditions favourable to the survival of GMOs (see Holmes, 1995 and for a review Doyle et al., 1995).

Moreover, there is no way to extrapolate from one region or environment to another, differing, environment. This is especially true when GMOs are transferred to ecosystems and climates which differ greatly from those where they were first developed and used. This is evident and acknowledged for deliberate release but also holds true for contained use in production plants with their multiple pathways for escape. Therefore, the scope of the biosafety protocol should cover the transfer of GMOs intended for contained use and have these included in the Advanced Informed Agreement procedures.

## References

- Aardema BW, Lorenz, MG and Krumbein WE (1983) Protection of sediment-adsorbed transforming DNA against enzymatic inactivation. *Applied and Environmental Microbiology* 46, 417-420.
- Adamo JA, Gealt MA (1996) A demonstration of bacterial conjugation within the alimentary canal of Rhabditis nematodes. *FEMS Microbiology Ecology* 20, 15-22
- Ahl T, Christoffersen K, Riemann B, Nybroe O (1995) A combined microcosm and mesocosm approach to examine factors affecting survival and mortality of *Pseudomonas fluorescens* Ag1 in seawater. *FEMS Microbiology Ecology* 17, 107-116
- Alvarez AJ, Yumet GM, Santiago CL, Toranzos GA (1996a) Stability of manipulated plasmid DNA in aquatic environments. *Environmental Toxicology and Water Quality* 11, 129-135
- Alvarez AJ, Yumet GM, Santiago CL, Hazen TC, Chaudhry R, Toranzos GA (1996b) In situ survival of genetically engineered microorganisms in a tropical aquatic environment. *Environmental Toxicology and Water Quality* 11, 21-25
- Amabile-Cuevas CF, Chicurel ME (1996) A possible role for plasmids in mediating the cell-cell proximity required for gene flux. *J. Theor. Biol.* 181, 237-243
- Angle JS, Levin MA, Gagliardi JV, McIntosh MS (1995) Validation of microcosms for examining the survival of *Pseudomonas aureofaciens* (lacZY) in soil. *Applied and Environmental Microbiology* 61, 2835-2839
- Armon R, Dosoretz C, Yoirish A, Shelef G, Neeman I (1995) Survival of the phytopathogen *Erwinia carotovora* subsp. *Carotovora* in sterile and non-sterile soil, sand and their admixture. *Journal of Applied Bacteriology* 79, 513-518
- Arturo-Schaan M, Tamanai-Shacoori Z, Thomas D, Cormier M (1996) Variations in R-plasmid DNA concentrations of *Escherichia coli* during starvation in sewage and brackish waters. *Journal of Applied Bacteriology* 80, 117-123
- Ashelford KE, Fry JC, Learner MA (1995) Plasmid transfer between strains of *Pseudomonas putida*, and their survival, within a pilot scale percolating-filter sewage treatment system. *FEMS Microbiology Ecology* 18, 15-25
- Awong J, Bitton, G and Chaudhry, G R (1990) Microcosm for assessing survival of genetically engineered microorganisms in aquatic envi-

- ronments. *Applied and Environmental Microbiology* 56, 977-983.
- Bale MJ, Day MJ, Fry MC (1988) Novel method for studying plasmid transfer in undisturbed river epilithon. *Applied and Environmental Microbiology* 54, 2756-2758
- Barkay T, Kroer N, Rasmussen LD, Sorensen SJ (1995) Conjugal transfer at natural population densities in a microcosm simulating an estuarine environment. *FEMS Microbiology Ecology* 16, 43-53
- Bauda P, Lallement C, Manem J (1995) Plasmid content evaluation of activated sludge. *Water Research* 29, 371-374
- Baur B, Hanselmann K, Schlimme W, Jenni B (1996) Genetic transformation in freshwater: *Escherichia coli* is able to develop natural competence. *Applied and Environmental Microbiology* 62, 3673-3678
- Beebe TJ (1993) Identification and analysis of nucleic acids in natural freshwaters. *Science of the Total Environment* 135, 123-129
- Bej AK, Steffan RJ, DiCesare J, Haff L, Atlas RM (1990) Detection of coliform bacteria in water by polymerase chain reaction and gene probes. *Applied and Environmental Microbiology* 56, 307-314
- Bergemann K, Noe W, Walz F (1994) Identifizierung und Messung von Parametern zur Beurteilung der potentiellen Risiken bei der Entsorgung von rekombinanten tierischen Zellkulturen im Produktionsmaßstab. In: *Biologische Sicherheit/Forschung Biotechnologie. BMFT Germany* (Ed.) p. 579-632.
- Bianchi A, Giuliano L (1996) Enumeration of viable bacteria in the marine pelagic environment. *Applied and Environmental Microbiology* 62, 174-177
- Binnerup SJ, Jensen DF, Thordal-Christensen H, Sorensen J (1993) Detection of viable, but non-culturable *Pseudomonas fluorescens* DF57 in soil using a microcolony epifluorescence technique. *FEMS Microbiology Ecology* 12, 97-105
- Bogosian G, Violand BN, Dorward-King EJ, Workman WE, Jung PE, Kane JF (1989) Biosynthesis and incorporation into protein of norleucine by *Escherichia coli*. *Journal of Biology and Chemistry* 264, 531-539.
- Boivin-Jahns V, Ruimy R, Bianchi A, Daumas S, Christen R (1996) Bacterial diversity in a deep-subsurface clay environment. *Applied and Environmental Microbiology* 62, 3405-3412
- Boyle-Vavra S, Seifert HS (1996) Uptake-sequence-independent DNA transformation exists in *Neisseria gonorrhoeae*. *Microbiology* 142, 2839-2845
- Brettar I, Ramos-Gonzalez MI, Ramos JL, Höfle MG (1994) Fate of



- Pseudomonas putida* after release into lake water mesocosms: different survival mechanisms in response to environmental conditions. *Microbial Ecology* 27, 99-122
- Brockmann E, Jacobsen BL, Hertel C, Ludwig W, Schleifer KH (1996) Monitoring of genetically modified *Lactococcus lactis* in gnotobiotic and conventional rats by using antibiotic resistance markers and specific probe or primer based methods. *Systematic and Applied Microbiology* 19, 203-212
- Byrd JJ, Colwell RR (1990) Maintenance of Plasmids pBR322 and pUC8 in nonculturable *Escherichia coli* in the marine environment. *Applied and Environmental Microbiology* 56, 2104-2107.
- Byrd JJ, Leahy JG, Colwell RR (1992) Determination of plasmid DNA concentration maintained by nonculturable *Escherichia coli* in marine microcosms. *Applied and Environmental Microbiology* 58, 2266-2270
- Byzov BA, Ngugen T, Babjeva IP (1993) Yeasts associated with soil invertebrates. *Biology and Fertility of Soils* 16, 183-187
- Canteras JC, Juanes JA, Perez L, Koev KN (1995) Modelling the coliforms inactivation rates in the Cantabrian Sea (Bay of Biscay) from in situ and laboratory determinations of T-90. *Water Science and Technology* 32, 37-44
- Chamier B, Lorenz MG, Wackernagel W (1993) Natural transformation of *Acinetobacter calcoaceticus* by plasmid DNA adsorbed on sand and groundwater aquifer material. *Applied and Environmental Microbiology* 59, 1662-1667.
- Cheng Q, Campbell EA, Naughton AM, Johnson S, Masure HR (1997) The *com* locus controls genetic transformation in *Streptococcus pneumoniae*. *Molecular Microbiology* 23, 683-692
- Christoffersen K, Ahl T, Nybroe O (1995) Grazing of nonindigenous bacteria by nano-sized protozoa in a natural coastal system. *Microbial Ecology* 30, 67-78
- Cirvilleri G, Marino R, Catara V, Di Silvestro S (1996) Monitoring of *Pseudomonas fluorescens* released in the phylloplane of different species in microcosm. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz* 103, 507-516
- Clegg CD, Anderson JM, Lappinscott HM, Van Elsas JD, Jolly J M (1995) Interaction of a genetically modified *pseudomonas fluorescens* with the soil-feeding earthworm *Octolasion cyaneum* (lumbricidae). *Soil Biology and Biochemistry* 27, 1423 -1429
- Clegg CD, Van Elsas JD, Anderson JM, Lappinscott HM (1996) Survival

- of parental and genetically modified derivatives of a soil isolated *Pseudomonas fluorescens* under nutrient-limiting conditions. *Journal of Applied Bacteriology* 81, 19-26
- Collard JM, Provoost A, Dijkmans R, Mergeay M (1993) Heavy metal-resistance genes as markers for measuring gene transfer in soils. In: *Bridge/Biotech, final sectorial meeting on biosafety and first sectorial meeting on microbial ecology*. Granada, October 24-27, 1993, p. 43
- Colwell RR, Brayton P, Herrington D, Tall B, Huq A, Levine MM (1996) Viable but non-culturable *Vibrio cholerae* 01 revert to a cultivable state in the human intestine. *World Journal of Microbiology and Biotechnology* 12, 28-31
- Daane LL, Molina JAE, Berry EC, Sadowsky MJ (1996) Influence of Earthworm Activity on Gene Transfer from *Pseudomonas fluorescens* to Indigenous Soil Bacteria. *Applied and Environmental Microbiology* 62, 515-521
- Dane F, Shaw JJ (1996) Survival and Persistence of Bioluminescent *Xanthomonas campestris* pv. *campestris* on Host and Non-Host Plants in the Field Environment. *Journal of Applied Bacteriology* 80, 73-80
- Davies CM, Long JAH, Donald M, Ashbolt NJ (1995) Survival of fecal microorganisms in marine and freshwater sediments. *Applied and Environmental Microbiology* 61, 1888-1896
- De Leij FAAM, Sutton EJ, Whipps JM, Fenlon JS, Lynch JM (1995) Impact of field release of genetically modified *Pseudomonas fluorescens* on indigenous microbial populations of wheat. *Applied and Environmental Microbiology* 61, 3443-3453
- De Rore H, Demolder K, De Wilde K, Top E, Houwen F, Verstraete W (1994a) Transfer of the catabolic plasmids RP4: Tn4271 to indigenous soil bacteria and its effect on respiration and biphenyl breakdown. *FEMS Microbiology Ecology* 15, 71-78
- De Rore H, Top E, Houwen F, Mergeay M, Verstraete W (1994b) Evolution of heavy metal resistant transconjugants in a soil environment with a concomitant selective pressure. *FEMS Microbiology Ecology* 14, 263-274
- Del Solar G, Moscoso M, Espinosa M (1993). Promiscuous plasmid from gram-positive bacteria as vehicles for horizontal gene transfer. In: *Bridge/Biotech; final sectorial meeting on biosafety and first sectorial meeting on microbial ecology*. Granada, October 24-27, 1993, p. 23
- Doerfler W, Schubbert R (1997) Fremde DNA I im Säugersystem. *Deutsches Ärzteblatt* 51/52 A 3465-3470

- Dott W, Khoury N, Ankel-Fuchs D, Henninger W, Kämpfer P (1991). Überlebensfähigkeit von genetisch veränderten *Escherichia coli*-Stämmen. 2. Mitteilung: Überleben von Reinkulturen in verschiedenen Wasser- und Bodenmatrices. *Zentralblatt Hygiene* 192: 1-13.
- Doucet-Populaire F (1992) Conjugal transfer of genetic information in gnotobiotic mice. In: *Microbial Releases*. Gauthier MJ (Ed.) Springer Verlag, Berlin, p. 161-166.
- Doyle JD, Stotzky G, McClung G, Hendricks CW (1995) Effects of genetically engineered micororganisms on microbial populations and processes in natural habitats. *Advances in Applied Microbiology* 40, 237-287
- Dresing U, Damman-Kalinowski T, Keller M, Selbitschka W, Pühler A, Tebbe C, Schwieger F, Munch JC (1995) Persistence of two bioluminescent *Rhizobium meliloti* strains in model ecosystems and in field release experiments. in: *Abstracts of the annual meeting of the genetic society 1995 in Bielefeld*. Kalinowski J, Pühler A, Schäfer A (Eds.). Gesellschaft für Genetik Köln, 18
- Dunn A, Day M, Randerson P (1993) Plasmid replication and maintenance in binary fissile microorganisms. *Endeavour* 17, 21-25.
- Duval-Ifлах Y (1992) Gene transfer in human and animals gut. In: *Microbial Releases*. Gauthier MJ (Ed.) Springer Verlag, Berlin, p. 151-160
- Edwards, C. 1993. The significance of in situ activity on the efficiency of monitoring methods, p.1-25. In: *Monitoring genetically manipulated microorganisms in the environment*. Edwards C (Ed.). John Wiley and Sons Ltd., New York.
- England LS, Lee H, Trevors JT (1995) Recombinant and wild-type *Pseudomonas aureofaciens* strains introduced into soil microcosms - effect on decomposition of cellulose and straw. *Molecular Ecology* 4, 221-230
- Feldmann SD, Sahm H (1994). Untersuchungen zum Verhalten gentechnisch veränderter Mikroorganismen in Laborkläranlagen. *BIOforum* 6, 220-226.
- Ferguson Y, Glover LA, McGillivray DM, Prosser JI (1995) Survival and activity of lux-marked *Aeromonas salmonicida* in seawater. *Applied and Environmental Microbiology* 61, 3494-3498
- Ferianc P, Polek B, Godocikova J, Toth D (1995) Metabolic activity of cadmium-stressed and /or starved *Vibrio* sp. *Folia Microbiologica* 40, 443-446
- Fish JT, Pettibone GW (1995) Influence of freshwater sediment on the sur-

- vival of *Escherichia coli* and *Salmonella* sp. as measured by three methods of enumeration. *Letters in Applied Microbiology* 20, 277-281
- Flint KP (1987). The long-term survival of *Escherichia coli* in river water. *Journal of Applied Bacteriology* 63, 261-270
- Frischer ME, Stewart GJ, Paul JH (1994) Plasmid transfer to indigenous marine bacterial populations by natural transformation. *FEMS Microbiology Ecology* 15, 127-135
- Frischer ME, Thurmond JM, Paul JH (1993) Factors affecting competence in a high frequency of transformation marine *Vibrio*. *Journal of General Microbiology* 139, 753-761
- Fry JC, Day MJ (1990) *Bacterial Genetics in Natural Environments*. Chapman and Hall, London, New York
- Fujimura H, Sakuma Y, Amann E (1994) Survival of genetically-engineered and wild-type strains of the Yeast *saccharomyces cerevisiae* under simulated environmental conditions - a contribution on risk assessment. *Journal of Applied Bacteriology* 77, 689-693
- Fulthorpe RA, Wyndham RC (1991) Transfer and expression of the catabolic plasmid pBRC60 in wild bacterial recipients in a freshwater ecosystem. *Applied and Environmental Microbiology* 57, 1546-1553
- Gadkari D (1991) Conjugation between two *Escherichia coli* strains and between *Escherichia coli* (donor) and *Azospirillum brasilense* (recipient) in the presence of various herbicides. *Toxicological and Environmental Chemistry* 30, 211-217
- Garcia-Lara J, Martinez J, Vilamu M, Vives-Rego J (1993) Effect of previous growth condition on the starvation-survival of *Escherichia coli* in seawater. *Journal of General Microbiology* 139, 1425-1431
- Gillespie KM, Angle JS, Hill RL (1995) Runoff losses of *Pseudomonas aureofaciens* (lacZY) from soil. *FEMS Microbiology Ecology* 17, 239-245
- Goodman AE, Marshall KC, Hermansson M (1994) Gene transfer among bacteria under conditions of nutrient depletion in simulated and natural aquatic environments. *FEMS Microbiology Ecology* 15, 55-60
- Goodwin D, Slater JH (1979) The influence of the growth environment on the stability of a drug resistance plasmid in *Escherichia coli* K12. *Journal of General Microbiology* 111, 201-210
- Götz A, Pukall R, Smit E, Tietze E, Prager R, Tschäpe H, Van Elsland JD, Smalla K (1996) Detection and characterization of broad-host-range plasmids in environmental bacteria by PCR. *Applied and Environmental*

*tal Microbiology* 62, 2621-2628

- Götz A, Smalla K (1997) Manure enhances plasmid mobilization and survival of *Pseudomonas putida* introduced into field soil. *Applied and Environmental Microbiology* 63, 1980 -1986
- Gross A, Wurz A, Willmund R (1994) Untersuchungen zum Verbleib rekombinanter Plasmid-DNA in einer Modellkläranlage. Korrespondenz Abwasser.
- Heijnen CE, Marinissen JCY (1995) Survival of bacteria introduced into soil by means of transport by *Lumbricus rubellus*. *Biology and Fertility of Soils* 20, 63-69
- Heinemann JA (1991) Genetics of gene transfer between species. *TIG* 7, 181-185
- Heinemann JA (1997) Assessing the risk of interkingdom DNA transfer, in: *Nordic seminar on antibiotic resistance marker genes and transgenic plants*. The Norwegian Biotechnology Advisory Board, 17-28
- Heitkamp MA, Kane JF, Morris PJL, Bianchini M, Hale MD, Bogosian G (1993) Fate in sewage of a recombinant *Escherichia coli* K-12 strain used in the commercial production of bovine somatotropin. *Journal of Industrial Microbiology* 11, 243-252
- Hekman WE, Heijnen CE, Burgers SLGE, VanVeen JA, Van Elsas JD (1995) Transport of bacterial inoculants through intact cores of two different soils as affected by water percolation and the presence of wheat plants. *FEMS Microbiology Ecology* 16, 143 -157
- Henschke RB, Schmidt FRJ (1990) Plasmid mobilization from genetically engineered bacteria to members of the indigenous soil microflora in situ. *Current Microbiology* 20, 105-110
- Hermansson M, Linberg C (1994) Gene transfer in the marine environment. *FEMS Microbiology Ecology* 15, 47-54
- Hobom B (1995) Kein Platz für verirrte Mikroben. Ergebnisse des Projektbeirates des Verbundprojektes Sicherheitsforschung Gentechnik
- Hodson RE, Dustman WA, Garg RP, Moran MA (1995) In situ PCR for visualization of microscale distribution of specific genes and gene products in prokaryotic communities. *Applied and Environmental Microbiology* 61, 4074-4082
- Holmes MT (1995) Ecological assessment after the addition of genetically engineered *Klebsiella planticola* SDF20 into soil, PhD-thesis, Oregon State University

- Hugenholz P, Pace NR (1996) Identifying microbial diversity in the natural environment: a molecular phylogenetic approach. *Trends in Biotech.* 14, 190-197
- Inamori Y; Murakami K, Sudo R, Kurihara Y, Tanaka N (1992) Environmental assessment method for field release of genetically engineered microorganisms using microcosm systems. *Wat. Sci. Tech.* 26, 2161-2164
- Istock CA (1991) Genetic exchange and genetic stability in bacterial populations. In: *Assessing ecological risks of biotechnology*. Ginzburg, R. (Ed.). Butterworth-Heinemann, Stoneham, MA, p.123-150.
- Jain RK, Sayler GS, Wilson JT, Houston L, Pacia D (1987) Maintenance and stability of introduced genotypes in groundwater aquifer material. *Applied and Environmental Microbiology* 53, 996-1002
- Jensen GB, Andrup L, Wilcks A, Smidt L, Poulsen OM (1996) The aggregation-mediated conjugation system of *Bacillus thuringiensis* subsp. *israelensis*: Host range and kinetics of transfer. *Current Microbiology* 33, 228-236
- Jiang SC, Paul JH (1995) Viral contribution to dissolved DNA in the marine environment as determined by differential centrifugation and Kingdom probing. *Applied and Environmental Microbiology* 61, 317-325
- Jiang XP, Chai TJ (1996) Survival of *Vibrio parahaemolyticus* at low temperatures under starvation conditions and subsequent resuscitation of viable, nonculturable cells. *Applied and Environmental Microbiology* 62, 1300-1305
- Jones N (1994) Relaxing European Regulations, "changes said to be necessary to take account of recognition that the risks to human and environmental safety are lower than was thought when the directives were adopted". *Bio/Technology* 12, 1144
- Karapinar M, Sahika AG (1991) Survival of *Yersinia enterocolitica* and *Escherichia coli* in spring water. *International Journal of Food Microbiology* 13, 315 -320
- Khanna M, Stotzky G (1992) Transformation of *Bacillus subtilis* by DNA bound on montmorillonite and effect of DNase on the transforming ability of bound DNA. *Applied and Environmental Microbiology* 58,1930-1939
- Kidambi SP, Ripp , Miller RV (1994) Evidence of phage-mediated gene transfer among *Pseudomonas aeruginosa* strains on the phylloplane. *Applied and Environmental Microbiology* 60, 496-500
- Kinkel LL, Wilson M, Lindow SE (1996) Utility of microcosm studies for

- predicting phylloplane bacterium population sizes in the field. *Applied and Environmental Microbiology* 62, 3413-3423
- Klier A (1992) Release of genetically modified microorganisms in natural environments: Scientific and ethical problems. In: *Microbial Releases*. Gauthier MJ (Ed.). Springer Verlag, Berlin, p. 183-190
- Klijn N, Weerkamp AH, De Vos WM (1995a) Detection and characterization of lactose-utilizing *Lactococcus* spp. in natural ecosystems. *Applied and Environmental Microbiology* 61, 788-792
- Klijn N, Weerkamp AH, De Vos WM (1995b) Genetic marking of *Lactococcus lactis* shows its survival in the human gastrointestinal tract. *Applied and Environmental Microbiology* 61, 2771-2774
- Klingmüller W (1993) Plasmid transfer in natural soil, as stimulated by sucrose, wheat rhizosphere, and ground sugar beets: a study with nitrogen-fixing enterobacter. *Microb. Releases* 1, 229-235
- Klingmüller W, Rieder G (1994) A) Konjugativer Transfer gentechnisch veränderter Plasmide B) Konjugativer Transfer gentechnisch veränderter Plasmide bei Freisetzungssimulation. In: *Biologische Sicherheit/Forschung Biotechnologie*. BMFT Germany (Ed.) Vol. 3, p. 83-106
- Kokjohn TA, Miller RV (1992) Gene transfer in the environment: transduction. In: *Release of genetically engineered and other micro-organisms*. Fry JC, Day M (Eds.) Cambridge University Press, 54-81
- Korhonen LK, Martikainen PJ (1991) Survival of *Escherichia coli* and *Campylobacter jejuni* in untreated and filtered lake water. *Journal of Applied Bacteriology* 71, 379-382
- Kozdroj J, Piotrowska-Seget Z (1995) Indigenous microflora and bean responses to introduction of genetically modified *Pseudomonas fluorescens* strains into soil contaminated with copper. *Journal of Environmental Science and Health Part A* 30, 2133-2158
- Kragelund L, Nybroe O (1996) Competition between *Pseudomonas fluorescens* Ag1 and *Alcaligenes eutrophus* JMP134 (pHP4) during colonization of barley roots. *FEMS Microbiology Ecology* 20, 41-51
- Kruse H, Sørum H (1994) Transfer of multiple drug resistance plasmids between bacteria of diverse origins in natural microenvironments. *Applied and Environmental Microbiology* 60, 4015-4021
- Lafuente R, Maymó-Gatell X, Mas-Castellà J, Guerrero R (1996) Influence of environmental factors on plasmid transfer in soil microcosms. *Current Microbiology* 32, 213-220
- Lebaron P, Batailler N, Baleux B (1994) Mobilization of a recombinant

- nonconjugative plasmid at the interface between wastewater and the marine coastal environment. *FEMS Microbiology Ecology* 15, 61-70
- Lee SY, Bollinger J, Bezdicek D, Ogram A (1996) Estimation of the abundance of an uncultured soil bacterial strain by a competitive quantitative PCR method. *Applied and Environmental Microbiology* 62, 3787-3793
- Leff LG, Leff AA (1996) Use of green fluorescent protein to monitor survival of genetically engineered bacteria in aquatic environments. *Applied and Environmental Microbiology* 62, 3486-3488
- Leser TD (1995) Validation of microbial community structure and ecological functional parameters in an aquatic microcosm designed for testing genetically engineered microorganisms. *Microbial Ecology* 29, 183-201
- Levy SB, Marshall B, Rowse-Eagle D (1980) Survival of *Escherichia coli* host-vector systems in the mammalian intestine. *Science* 209, 391-394
- Lindahl V (1996) Improved soil dispersion procedures for total bacterial counts, extraction of indigenous bacteria and cell survival. *Journal of Microbiological Methods* 25, 279-286
- Lindahl V, Bakken LR (1995) Evaluation of methods for extraction of bacteria from soil. *FEMS Microbiology Ecology* 16, 135-142
- Lorenz MG, Aardema BW, Wackernagel W (1988) Highly efficient genetic transformation of *Bacillus subtilis* attached to sand grains. *Journal of General Microbiology* 134, 107-112
- Lorenz MG, Wackernagel W (1987) Adsorption of DNA to sand and variable degradation rates of adsorbed DNA. *Applied and Environmental Microbiology* 53, 2948-2952
- Lorenz MG, Wackernagel W (1992) DNA binding to various clay minerals and retarded enzymatic degradation of DNA in a sand/clay microcosm. In: *Microbial Releases*. Gauthier MJ (Ed.). Springer Verlag, Berlin, p. 103-113
- Lorenz MG, Wackernagel W (1993) Bacterial gene transfer in the environment. In: *Transgenic organisms*. Wöhrmann K, Tomiuk J (Eds.) Birkhäuser, Basel, Boston, Berlin, p 43-64
- Lorenz MG, Wackernagel W (1994) Bacterial gene transfer by natural genetic transformation in the environment. *Microbiological Reviews* 58, 563-602
- Lou WQ, Yousef AE (1996) Resistance of *Listeria monocytogenes* to heat after adaptation to environmental stresses. *Journal of Food Protection* 59, 465-471



- Lukin SA, Prozorov AA (1992) Conjugate transfer of plasmid dna between bacteria in soil. *Microbiology* 61, 493-496
- Lunsford RD, London J (1996) Natural genetic transformation in *Streptococcus gordonii*: comX imparts spontaneous competence on strain wicky. *Journal of Bacteriology* 178, 5831-5835
- Mazodier P, Davies J (1991) Gene transfer between distantly related bacteria. *Annual Reviews on Genetics* 25, 147-171
- McClure NC, Weightman AJ, Fry JC (1989) Survival of *Pseudomonas putida* UWC1 containing cloned catabolic genes in a model activated-sludge unit. *Applied and Environmental Microbiology* 55, 2627-2634
- Mezrioui N, Baleux B, Troussellier M (1995) A microcosm study of the survival of *Escherichia coli* and *Salmonella typhimurium* in brackish water. *Water Research* 29, 459 -465
- Mieschendahl M, Danneberg G (1994) Untersuchungen zur in vivo-Transformation von Mikroorganismen in Oberflächengewässern. In: *Biologische Sicherheit/Forschung Biotechnologie*. BMFT Germany (Ed.). Vol. 3, p. 901-916
- Miller RV, Ripp S, Replicon J, Ogunseitan OA, Kokjohn TA (1992) Virus-mediated gene transfer in freshwater environments. In: *Microbial Releases*. Gauthier MJ (Ed.). Springer Verlag, Berlin, p. 51-62
- Morrison M (1996) Do ruminal bacteria exchange genetic material? *J. Dairy Sci.* 79, 1476-1486
- Naik GA, Bhat, LN, Chopade, BA, Lynch, JM (1994). Transfer of broad-host range antibiotic resistance plasmids in soil microcosms. *Current Microbiology* 28, 209-215
- Nwoguh CE, Harwood CR, Barer MR (1995) Detection of induced beta-galactosidase activity in individual non-culturable cells of pathogenic bacteria by quantitative cytological assay. *Molecular Microbiology* 17, 545-554
- Nybroe O, Einarson K, Ahl T (1996) Growth and viability of *Alcaligenes eutrophus* JMP134 in seawater as affected by substrate and nutrient amendment. *Letters in Applied Microbiology* 22, 366-370
- Ogunseitan OA, Salyer GS, Miller RV (1992) Application of DNA probes to the analysis of bacteriophage distribution in the environment. *Applied and Environmental Microbiology* 58, 2046-2052
- Oliver JD, Hite F, McDougald D, Andon NL, Simpson LM (1995a) Entry into, and resuscitation from, the viable but nonculturable state by *Vibrio vulnificus* in an estuarine environment. *Applied and Environmental Microbiology* 61, 2624-2630

- Oliver JD, McDougald D, Barrett T, Glover LA, Prosser JI (1995b) Effect of temperature and plasmid carriage on nonculturability in organisms targeted for release. *FEMS Microbiology Ecology* 17, 229-237
- Overbeck J (1991) Ermittlung und Bewertung des ökologischen Risikos im Umgang mit gentechnisch veränderten Organismen. Vortrag und Diskussion zum Vortrag, Forschungsbericht: 10802 086/01. Umweltbundesamt.
- Pace NR (1997) A molecular view of microbial diversity and the biosphere. *Science* 276, 734-730
- Paget E, Jouan D, Lebrun M, Simonet P (1993) Persistence of plant DNA in soil. In: *Bridge/Biotech; final sectorial meeting on biosafety and first sectorial meeting on microbial ecology*, Granada, October 24-27, 1993, p. 33
- Paget E, Monrozier LJ, Simonet P (1992) Adsorption of DNA on clay minerals: protection against DNase I and influence on gene transfer. *FEMS Microbiology Letters* 97, 31-40
- Paget E, Simonet P (1993) Evidence of gene transfer in soil via transformation. In: *Bridge/Biotech; Final sectorial meeting on biosafety and first sectorial meeting on microbial ecology*, Granada, October 24-27, 32
- Paget E, Simonet P (1994) On the track of natural transformation in soil. *FEMS Microbiology Ecology* 15, 109-118
- Paul JH (1992) Intergeneric natural plasmid transformation between *Escherichia coli* and a marine *Vibrio* species. In: *Microbial Releases*. Gauthier; M.J. (Ed.). Springer Verlag, Berlin, p. 63-68
- Paul JH, Frischer ME, Thurmond JM (1991) Gene transfer in marine water column and sediment microcosms by natural plasmid transformation. *Applied and Environmental Microbiology* 57 1509-1515
- Pickup RW (1991) Development of molecular methods for the detection of specific bacteria in the environment. *Journal of General Microbiology* 137, 1009-1019
- Pickup RW, Morgan JAW, Winstanley C (1993) In situ detection of plasmid transfer in the aquatic environment. In: *Monitoring genetically manipulated microorganisms in the environment*. Edwards C. (Ed.) John Wiley and Sons Ltd., Chichester, S. 61-82
- Prozorov AA (1997) Cell differentiation related to genetic transformation in bacteria and its regulation. *Microbiology* 66, 1-7
- Pukall R, Tschäpe H, Smalla K (1996) Monitoring the spread of broad host and narrow host range plasmids in soil microcosms. *FEMS Microbiology Ecology* 20, 53-66
- Rahman I, Shahamat M, Chowdhury MAR, Colwell RR (1996) Potential

- virulence of viable but nonculturable *Shigella dysenteriae* type 1. *Applied and Environmental Microbiology* 62, 115-120
- Ramos JL, Diaz E, Dowling D, DeLorenzo V, Molin S, O'Gara F, Ramos C, Timmis KN (1994) The behavior of bacteria designed for biodegradation (Review). *Bio/Technology* 12, 1349-1356
- Rang CU, Kennan RM, Midtvedt T, Chao L, Conway PL (1996) Transfer of the plasmid RP1 in vivo in germ free mice and in vitro in gut extracts and laboratory media. *FEMS Microbiology Ecology* 19, 133-140
- Recorbet G, Richaume A, Jocteur-Monrozier L (1995) Distribution of a genetically-engineered *Escherichia coli* population introduced into soil. *Letters in Applied Microbiology* 21, 38-40
- Recorbet G, Steinberg C, Faurie G (1992) Survival in soil of genetically engineered *Escherichia coli* as related to inoculum density, predation and competition. *FEMS Microbiology Ecology* 101, 251-260
- Replicon J, Frankfater A, Miller RV (1995) A continuous culture model to examine factors that affect transduction among *Pseudomonas aeruginosa* strains in freshwater environments. *Applied and Environmental Microbiology* 61, 3359-3366
- Ripp S, Miller RV (1995) Effect of suspended particulates on the frequency of transduction among *Pseudomonas aeruginosa* in a freshwater environment. *Applied and Environmental Microbiology* 61, 1214-1219
- Ripp S, Ogunseitian OA, Miller RV (1994) Transduction of a freshwater microbial community by a new *Pseudomonas aeruginosa* generalized transducing phage UT1. *Molecular Ecology* 3, 121-126
- Rollinger Y, Dott W (1987) Survival of selected bacterial strains in sterilized activated carbon filters and biological activated carbon units. *Applied and Environmental Microbiology* 53, 777-781
- Romanowski G, Lorenz MG, Sayler G, Wackernagel W (1992) Persistence of free plasmid DNA in soil monitored by various methods, including a transformation assay. *Applied and Environmental Microbiology* 58, S.3012-3019
- Romanowski G, Lorenz MG, Wackernagel W (1990) Bakterieller Gentransfer durch freie DNA im natürlichen Lebensraum: Studien an einem Bodenmodell. In: *Biologische Sicherheit/Forschung Biotechnologie*. Band 2, BMFT (Hrsg.) S. 11-26
- Romanowski G, Lorenz MG, Wackernagel W (1993a) Plasmid DNA in a groundwater aquifer microcosm - adsorption, DNase resistance and natural genetic transformation of *Bacillus subtilis*. — *Molecular Ecology* 2, 171-181

- Romanowski G, Lorenz MG, Wackernagel W (1993b) Use of polymerase chain reaction and electroporation of *Escherichia coli* to monitor the persistence of extracellular plasmid DNA introduced into natural soils. *Applied and Environmental Microbiology* 59, 3438-3446
- Rybachenko NE, Stepanova TV, Rumer LM, Mozgovaya IN, Byzov BA, Zvyagintsev DG (1996) Conjugal transfer of plasmid pAM beta 1 from *Streptococci* to bacilli in vermicompost. *Microbiology* 65, 663-667
- Sandaa RA, Enger O (1994) Transfer in marine sediments of the naturally occurring plasmid pRAS1 encoding multiple antibiotic resistance. *Applied and Environmental Microbiology* 60, 4234-4238
- Saunders JR, Saunders VA (1993) Genotypic and phenotypic methods for the detection of specific released microorganisms. In: *Monitoring genetically manipulated microorganisms in the environment*. Edwards C (Ed.). John Wiley and Sons Ltd., New York, p.27-59.
- Saye DJ, Miller RV (1989) The aquatic environment: consideration of horizontal gene transmission in a diversified habitat. In: *Gene transfer in the environment*. Levy SB, Miller RV (Eds.). McGraw-Hill, New York., p.223 ff
- Schäfer A (1996) Horizontaler Gentransfer - Mechanismen und biologische Sicherheit. *Biospektrum* 6.96, 23-29
- Schäfer A, Kalinowski J, Pühler A (1994) Increased fertility of *Corynebacterium glutamicum* recipients in intergeneric matings with *Escherichia coli* after stress exposure. *Applied and Environmental Microbiology* 60, 756-759.
- Schicklmaier P, Schmieger H (1995) Frequency of generalized transducing phages in natural isolates of the *Salmonella typhimurium* complex. *Applied and Environmental Microbiology* 61, 1637 -1640
- Schlüter K, Potrykus I (1996) Horizontaler Gentransfer von Transgenen Mikroorganismen (Bakterien und Pilzen) und seine ökologische Relevanz. In: *Gentechnisch veränderte krankheits- und schädlingsresistente Nutzpflanzen. Eine Option für die Landwirtschaft?* Schulte E, Käppeli O (Hrsg.) Band I. Materialien. Eine Publikation des Schwerpunktprogrammes Biotechnologie des Schweizerischen Nationalfonds, Bern.
- Schmidt F, Brokamp A, Henschke R, Henschke E, Happe B (1994) Möglichkeiten und Risiken der landwirtschaftlichen Nutzung gentechnologisch veränderter Mikroorganismen. In: *Biologische Sicherheit/Forschung Biotechnologie*. BMFT Germany (Ed.) Vol. 3, p. 985-1008

- Schmidt FRJ (1991) Die Bewertung von Daten zu den Risiken bei der Freisetzung von gentechnisch veränderten Bodenmikroorganismen. *BIOforum*, 9/91, 312-316
- Schneider E, Schaefer B, Feuerpfeil I (1994) Konjugativer Gentransfer und Überleben gentechnisch veränderter Bakterien in Abwasser und Oberflächenwasser unter simulierten Freisetzungsbedingungen. In: *Biologische Sicherheit/Forschung Biotechnologie*. Band 3, BMFT (Hrsg.) p. 107-130
- Schubbert R, Doerfler W, (1996) Food ingested foreign DNA is taken up via gastrointestinal wall epithelia and peyer's patches into peripheral leucocytes, spleen and liver. In: *Transgenic Organisms and Biosafety*. Schmidt ER, Hankeln Th. (Eds.) Springer Verlag Berlin, Heidelberg, New York, p. 132
- Selbitschka W, Pühler A (1994) Nachweis von Gentransfer in Umweltmedien. In: *Überwachungsmethoden Gentechnik - Nachweisverfahren für Mikroorganismen, Viren und Gene in der Umwelt*. Forschungsbericht 108 02 089/01. Umweltbundesamt (Hrsg.) Berlin
- Sjogren RE (1995) Thirteen-year survival study of an environmental *Escherichia coli* in field. *Water, Air and Soil Pollution* 81, 315-335
- Smalla K (1995) Horizontal gene transfer from transgenic plants into plant associated bacteria and soil bacteria. in: *Safety of transgenic crops. Environmental and agricultural considerations. Proceedings of the Basel Forum on Biosafety*, 2/1995
- Smith HW (1975) Survival of orally administered *E. coli* K12 in alimentary tract of man. *Nature* 255, 500-502
- Sobecky PA, Schell MA, Moran MA, Hodson RE (1992) Adaption of model genetically engineered microorganisms to lake water: growth rate enhancements and plasmid loss. *Applied and Environmental Microbiology* 58, 3630-3637
- Sobecky PA, Schell MA, Moran MA, Hodson RE (1996) Impact of a genetically engineered bacterium with enhanced alkaline phosphatase activity on marine phytoplankton communities. *Applied and Environmental Microbiology* 62, p. 6-12
- Spielman A, Regal P, Mundt C, Klinger T, Kapuscinski AR, Istock C, Ingham ER, Holmes M, Fagan, JB (1996) *Draft — Assessment of genetically Engineered Organisms in the Environment: The Puget Sound Workshop Biosafety Handbook*, Edmonds Institute
- Stotzky G, Babich H (1994) Fate of genetically-engineered microbes in

- natural environments. *Recombinant DNA Technical Bulletin* 7, 163-188
- Stotzky G, Zeph LR, Devanas MA (1991) Factors affecting the transfer of genetic information among microorganisms in soil. In: *Assessing ecological risks of biotechnology*; Ginzburg R (Ed.). Butterworth-Heinemann, Stoneham, MA
- Tamanai-Shacoori Z, Jolivet-Gougeon A, Cormier M (1996) Comparison of direct PCR and PCR amplification after DNA extraction for the detection of viable enterotoxigenic *Escherichia coli* in laboratory microcosms. *Journal of Microbiological Methods* 26, 21-26
- Tanaka N, Inamori Y, Murakami K, Akamatsu T, Kurihara Y (1994) Effect of species composition on stability and reproductivity of a small-scale microcosm system. *Water Science and Technology* 30, 125-131
- Tebbe CC, Vahjen W, Munch JC, Feldmann SD, Ney U, Sahn H, Gelissen G, Amore R, Hollenberg CP (1994a) Überleben der Untersuchungsstämme und Persistenz ihrer rekombinanten DNA. *BioEngineering* 6/94, 14-21
- Tebbe CC, Vahjen W, Munch J, Meier B, Gellissen G, Feldmann S, Sahn H, Amore R, Hollenberg CP, Blum S, Wackernagel W (1994b) Mesokosmenuntersuchungen und Einfluß der Habitatbedingungen auf die Expression, Überdauerung und Übertragung des Aprotinins. *BioEngineering* 6/94, 22-26
- Temann UA, Hösl R, Klingmüller W (1992) Plasmid transfer between *Pseudomonas aeruginosa* strains in natural soil. In: *The release of genetically modified microorganisms*, Stewart-Tull DES, Sussmann M (Eds.), Plenum Press, New York
- Terzieva SI, McFeters GA (1991) Survival and injury of *Escherichia coli*, *Campylobacter jejuni*, and *Yersinia enterocolitica* in stream water. *Can. J. Microbiology* 37, 785-790
- Thompson IP, Ellis RJ, Bailey MJ (1995a) Autecology of a genetically modified fluorescent *Pseudomonad* on sugar beet. *FEMS Microbiology Ecology* 17, 1-13
- Thompson IP, Lilley AK, Ellis RJ, Bramwell PA, Bailey MJ (1995b) Survival, colonization and dispersal of genetically modified *Pseudomonas fluorescens* SBW25 in the phytosphere of field grown sugar beet. *Bio/Technology* 13, 1493-1497
- Tokuda Y, Ano T, Shoda M (1995) Survival of *Bacillus subtilis* NB22 and its transformant in soil. *Applied Soil Ecology* 2, 85-94
- Trieu-Cuot P, Carlier C, Martin P, Courvalin P (1987) Plasmid transfer by

- conjugation from *Escherichia coli* to gram-positive bacteria. *FEMS Microbiology Letters* 48, 289-294
- Tsushima S, Hasebe A, Komoto Y, Carter JP, Miyashita K, Yokoyama K, Pickup RW (1995) Detection of genetically engineered microorganisms in paddy soil using a simple and rapid nested polymerase chain reaction method. *Soil Biology and Biochemistry* 27, 219-227
- Turpin PE, Maycroft KA, Rowlands CL, Wellington EMH (1993) Viable but non-culturable salmonellas in soil. *Journal of Applied Bacteriology* 74, 421-427
- Vahjen W, Tebbe CC (1994) Enhanced detection of genetically engineered *Corynebacterium glutamicum* pUN1 in directly extracted DNA from soil, using the T4 gene 32 protein in the polymerase chain reaction. *Eur. J. Soil Biol.* 30, 93 -98
- Van Elsas JD (1992). Environmental pressure imposed on GEMMOS in soil. In: *The release of Genetically Modified Microorganisms*. Stewart-Tull DES, Sussman M (Eds.), Plenum Press / New York, 1-14
- Van Elsas JD, Trevors JT, Starodub ME, van Overbeek LS (1990) Transfer of plasmid RP4 between *Pseudomonas* after introduction into soil; influence of spatial and temporal aspects of inoculation. *FEMS Microbiology Ecology* 73, 1-12
- Van Overbeek LS, Eberl L, Givskov M, Molin S, Van Elsas JD (1995) Survival of, and induced stress resistance in, carbon-starved *Pseudomonas fluorescens* cells residing in soil. *Applied and Environmental Microbiology* 61, 4202-4208
- Van Der Hoeven N, Van Elsas JD, Heijnen CE (1996) A model based on soil structural aspects describing the fate of genetically modified bacteria in soil. *Ecological Modelling* 89, 161 -173
- Wackernagel W, Romanowski G, Lorenz MG (1992) Studies on gene flux by free bacterial DNA in soil, sediment and groundwater aquifer. In: *The release of genetically modified microorganisms*, Stewart-Tull DES, Sussman M, Plenum Press, New York, 171-173
- Wagner M, Amann R, Lemmer H, Schleifer KH (1993) Probing activated sludge with oligonucleotides specific for proteobacteria: inadequacy of culture-dependent methods for describing microbial community structure. *Applied and Environmental Microbiology* 59, 1520-1525
- Weinbauer MG, Fuks D, Puskaric S, Peduzzi P (1995) Diel, seasonal, and depth-related variability of viruses and dissolved DNA in the northern adriatic sea. *Microbial Ecology* 30, 25-41
- Weiskel PK, Howes BL, Heufelder GR (1996) Coliform contamination of

- a coastal embayment: Sources and transport pathways. *Environmental Science and Technology* 30, 1872-1881
- Wendt-Potthoff K, Backhaus H, Smalla K (1994) Monitoring the fate of genetically engineered bacteria sprayed on the phylloplane of bush beans and grass. *FEMS Microbiology Ecology* 15, 279-290
- Williams HG, Day MJ, Fry JC, Stewart GJ (1996) Natural transformation in river epilithon. *Applied and Environmental Microbiology* 62, 2994 - 2998
- Winkler J, Timmis KN, Snyder RA (1995) Tracking the response of *Burkholderia cepacia* G4 5223-PR1 in aquifer microcosms. *Applied and Environmental Microbiology* 61, 448 -455
- Wright DA, Killham K, Glover LA, Prosser JI (1995) Role of pore size location in determining bacterial activity during predation by protozoa in soil. *Applied and Environmental Microbiology* 61, 3537 -3543
- Yancey JrRJ, Kotarski SF, Thurn KK, Lepley RA, Mott JE (1993) Absence of persistence and transfer of genetic material by recombinant *Escherichia coli* in conventional, antibiotic-treated mice. *Journal of Industrial Microbiology* 11, 259-271
- Zweifel UL, Hagstrom A (1995) Total counts of marine bacteria include a large fraction of non-nucleoid-containing bacteria (ghosts). *Applied and Environmental Microbiology* 61, 2180-2185



## ABOUT THE AUTHORS

**Dr Beatrix Tappeser** is a biologist and coordinator of the Department of Risk Assessment in Genetic Engineering at the Institute for Applied Ecology, Freiburg, Germany. She has been working for fifteen years on questions concerning the possible social, ecological and health-related impacts of genetic engineering in different application fields. The Institute for Applied Ecology is a non-profit, citizen-founded research institution.

**Dr Manuela Jäger** studied biology at the University of Ulm, Germany, from 1983 to 1992, with microbiology, molecular genetics and virology as her main subjects. From 1992 to 1998, she was working at the Department of Risk Assessment in Genetic Engineering at the Institute for Applied Ecology, focusing on genetically engineered and novel foods and their impacts on human health.

**Dr Claudia Eckelkamp** is a biologist and has been a scientific collaborator at the Department of Risk Assessment in Genetic Engineering at the Institute for Applied Ecology since 1995. Her field of work covers risk assessment and evaluation of the exploitation of genetic engineering in different areas of application, particularly risk assessment of genetically modified microorganisms, environmental risks of the cultivation of genetically engineered plants in agriculture, and evaluation of the safety of genetically engineered foodstuffs.